

# Two Staining Methods in the Study of Ciliates

Xianyu Yang<sup>1,2,3\*</sup>, Xin Gao<sup>1,2,3</sup>

<sup>1</sup>College of Animal Science and Technology, Zhejiang A&F University, Hangzhou, China

<sup>2</sup>Key Laboratory of Applied Technology on Green-Eco-Healthy Animal Husbandry of Zhejiang Province, Hangzhou, China

<sup>3</sup>Zhejiang Provincial Engineering Laboratory for Animal Health Inspection and Internet Technology, Hangzhou, China

\*Corresponding author, email: yangxy78@zafu.edu.cn

---

**Abstract** — Ciliates have two kinds of nuclei, somatic macronuclei and germinal micronuclei. For the study of nuclear behaviour during their conjugation (sexual reproduction mode), nuclear staining techniques are very important. In the last decade, we studied nuclear behaviour of several species of *Paramecium*, and several nuclear staining methods were used. Here carbol fuchsin and “apofluor” (a double vital fluorescent dyes of acridine orange and Hoechst 33342) will be introduced.

**Keywords** — Ciliates; *Paramecium*; Carbol fuchsin; Acridine orange; Hoechst 33342

---

## I. INTRODUCTION

*Paramecium caudatum* is the representative of ciliates. As all other ciliates, it has two types of nuclei, a germinal micronucleus and a somatic macronucleus<sup>[1]</sup>. During the last decade, some new findings of nuclear behaviour during conjugation of several species of *Paramecium* were reported<sup>[2-8]</sup>, all of which benefited from the improvement of several experimental techniques<sup>[9-12]</sup> and adopting several staining methods, such as protargol<sup>[13,14]</sup>, immunofluorescence<sup>[15,16]</sup>, carbol fuchsin<sup>[17,18]</sup> and “apofluor”<sup>[19,20]</sup>. Recently, we have introduced these methods in detail. One is volume-fixing method suitable for the observation of living cells and photographing<sup>[21]</sup>. The second is a method how to collect large number of synchronized conjugating pairs<sup>[22]</sup>, and the third is a technique to get large amount of magnetic nanoparticles by ultrasonic treatment<sup>[23]</sup>. Here, we will mainly introduce two convenient staining methods, carbol fuchsin and “apofluor”.

## II. MATERIALS AND METHODS

### Main Equipments and Materials

Fluorescent microscope (Nikon 50i), acridine orange (AO, Shanghai Chemical Reagent Company, China), Hoechst 33342 (HO, Beyotime Institute of Biotechnology, China), carbol fuchsin solution (self-made), the exconjugant suspension of *P. caudatum*.

### Methods

#### Carbol fuchsin staining

According to Carr & Walker<sup>[17]</sup>, the protocol for the preparation of carbol fuchsin solution was summarized as following, which is scale down for its easy preparation. 1. Weigh 30 mg basic fuchsin and dissolve in 1 mL of 75% ethanol to make it completely dissolved. 2. Add 9 mL of 5% phenol aqueous solution to the above solution and mix well. 3. Add 1.08 mL acetic acid and formalin each into the above solution, respectively, and mix well. 4. Add 28 mL of 45% acetic acid aqueous solution into the above solution and mix well. 5. Finally, 40 mL carbol fuchsin staining solution would be obtained, which can be stored at room temperature for at least one year. 6. Before cell staining, the staining solution could be diluted with 45% acetic acid aqueous solution properly. 7. A temporary slide is made according to volume-fixing method (usually half of the cells moving slowly)<sup>[10,21]</sup>, then the staining solution is added on one side of the cover glass, and a piece of absorbent paper is used to suck the staining solution on the other side making the staining solution spread over the whole cover glass. 8. After 3-5 min, cells can be observed under the bright field microscope and photographed.

#### “Apofluor” staining

“Apofluor” is the combination of two vital fluorescent dyes, AO and HO<sup>[19]</sup>. The stock solutions (50x) of AO (1 µg/mL) and HO (10 µg/mL) prepared with autoclaved H<sub>2</sub>O should be kept at 4°C<sup>[20]</sup>. When it is ready for staining, 10 µl each is added to 480 µl exconjugant suspension of *P. caudatum*. About 10 min later, temporary

preparations are made by volume-fixing method<sup>[10,21]</sup>. Usually, 13  $\mu\text{L}$  cell suspension is corresponding to the cover glass of 22 mm\*22 mm, when about 50% of cells move very slowly being convenient for observation and photographing under Nikon 50i fluorescent microscopes with UV filters.

### III. RESULTS AND DISCUSSIONS

Concerning carbol fuchsin solution, nuclei on all developing stages are stained the same colour (Figure 1A-C). As to the "apofluor", contrast to the normal developing nuclei stained blue (Figure 1A'-C'), the apoptotic nuclei are stained yellow green (red arrows in Figure 1A').




Fig. 1 Exconjugants of *P. caudatum* stained by carbol fuchsin or "apofluor"  
A-C: Carbol fuchsin staining; A'-C': "Apofluor" staining. A, A': Cells on the metaphase of the third postzygotic division;  
B, B': Cells on the anaphase of the third postzygotic division; C, C': Cells on the telophase of the third postzygotic division.  
Red arrows: Apoptotic meiotic nuclei; Blue/white arrows: Non-degenerating normal nuclei. Bars: 20  $\mu\text{m}$ .

These above two staining methods have one common feature: speedy. Therefore, they are easy to be used. Of course, fluorescent microscope is indispensable for "apofluor" staining. In this sense, carbol fuchsin staining is the most convenient method, which is applicable to both the research and the student experiment of *P. caudatum*. However, "apofluor" is able to distinguish the apoptotic nuclei from normal/developing nuclei, which is convenient for studying apoptosis<sup>[2,20]</sup>.

In fact, we have tried these two staining methods to other ciliates, both work well<sup>[24]</sup>. Concerning protargol<sup>[13,14]</sup> and immunofluorescence<sup>[15,16]</sup>, they have their own advantages. Unique nuclear division of the third prezygotic division was indicated by protargol in *P. caudatum*<sup>[3]</sup> and *P. polycaryum*<sup>[14]</sup>. Cytoplasmic microtubules surrounding the surviving meiotic nuclei and the nuclei on-going the third postzygotic division were indicated by the immunofluorescence with anti-alpha tubulin antibody<sup>[7]</sup>. However, experiments with protargol and immunofluorescence could not always get ideal results.

#### ACKNOWLEDGMENT

This work was partially supported by the grants from National Natural Science Foundation of China (No. 31772409, 31372149).

## REFERENCES

- [1] Wichterman R. *The Biology of Paramecium*, 2nd ed. Plenum Press, New York and London, 1986.
- [2] Gao X, Zhang X, Yang X. Morphological apoptotic characteristics of the post-meiotic micronuclei in *Paramecium caudatum*. *Eur J Protistol*, 2010, 46(3): 243-250.
- [3] Gao X, Shi X, Yang X. Dynamic behaviour of stationary pronuclei during their positioning in *Paramecium caudatum*. *Eur J Protistol*, 2011, 47(3): 235-237.
- [4] Gao X, Xu C, Yang X. Characterization of meiotic nuclear degeneration in *Paramecium tetraurelia*. *Chn J Zool*, 2011, 46(3): 89-92. (in Chinese with English abstract)
- [5] Gao X, Zhu J, Yang X, Yuan J, Wang Y, Song M. Localization of stationary pronuclei during the conjugation of *Paramecium* indicated by immunofluorescence staining. *Zool Res*, 2011, 32(4): 461-464.
- [6] Gao X, Yang X, Zhu J, Yuan J, Wang Y, Song M. New details indicated by different stainings during the conjugation of ciliated protozoa *Paramecium*. *Zool Res*, 2011, 32(6): 651-656.
- [7] Wang Y, Yuan J, Gao X, Yang X. Stage-specific appearance of cytoplasmic microtubules around the surviving nuclei during the third prezygotic division of *Paramecium*. *Zool Res*, 2012, 33(E5-6): E97-E102.
- [8] Gao X, Song Y, Yang X. Nuclear behavior in exconjugants of *Paramecium bursaria*. *IJRDO-Journal Biological Science*, 2018, 4(7): 1-5.
- [9] Yang X, Takahashi M. Disturbance of the determination of germinal and somatic nuclei by heat shock in *Paramecium caudatum*. *J Eukaryot Microbiol*, 1999, 46(1): 49-55.
- [10] Lin Y, Gao X, Yang X. A method for observation of living *Paramecium*. *Shichuan J Zool*, 2009, 28: 276. (in Chinese with English abstract)
- [11] Vosskühler C, Tiedtke A. Magnetic separation of phagosomes of defined age from *Tetrahymena thermophila*. *J Eukaryot Microbiol*, 1993, 40: 556-562.
- [12] Sun X, Wu J, Gao X, Yang X. To enhance the production of magnetic iron-dextran particles by ultrasonic treatment and its applications in ciliate studies. *Chn J Zool*, 2010, 45(1): 90-93. (in Chinese with English abstract)
- [13] Shi X, Frankel J. Morphology and development of mirror-image doublets of *Stylonychia mytilus*. *J Protozool*, 1991, 37(1): 1-13.
- [14] Yang X, Shi X. Gametic nuclear exchange during the conjugation of *Paramecium polycaryum*. *JPN J Protozool*, 2007, 40: 113-121.
- [15] Ishida M, Nakajima Y, Kurokawa K, Mikami K. Nuclear behavior and differentiation in *Paramecium caudatum* analyzed by immunofluorescence with anti-tubulin antibody. *Zoo Sci*, 1999, 16: 915-926.
- [16] Yang X, Takahashi M. Nuclei may anchor at specific locations during nuclear determination in *Paramecium caudatum*. *Eur J Protistol*, 2002, 38(2): 147-153.
- [17] Carr DH, Walker JE. Carbol fuchsin as a stain for human chromosome. *Biotech Histochem*, 1961, 36(4): 233-236.
- [18] Yang X, Takahashi M. Stage-related profiles of the reaction of the cells to heat shock during meiotic prophase in *Paramecium caudatum*. *J Exp Zool*, 1996, 275: 53-60.
- [19] Santos ML, Lu E, Wolfe J. Nuclear death in living *Tetrahymena*: the case of the haploid nuclei. *J Eukaryot Microbiol*, 2000, 47: 493-498.
- [20] Yang X, Gao X, Shi X. Detection of haploid nuclear death in living *Paramecium caudatum*. *JPN J Protozool*. 2007, 40: 123-130.
- [21] Yang X, Gao X. Introduction of an efficient method for the observation of living *Paramecium*. *IJRDO-Journal of Education Research*, 2018, 3(10): 27-29.
- [22] Gao X, Yang X. Introduction of an effective method for separating conjugating pairs of *Paramecium*. *IJRDO-Journal of Biological Science*, 2019, 5(10): 1-3.
- [23] Gao X, Yang X. Increasing the yield of iron-dextran nanoparticles by ultrasonic treatment. *IJRDO-Journal of Biological Science*, 2019, 5(10): 4-6.
- [24] Yang X, Zhang X, Huang X. The application of vital fluorescence dyes in the study of ciliates. *Acta Hydrobiol Sinica*, 2008, 32(suppl): 80-83. (in Chinese with English abstract)